



Effective Viscosity of Conned Hydrocarbons

Sivebæk, Ion Marius; Samoilov, Vladimir N.; Persson, Bo N.J.

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Sivebæk, I. M., Samoilov, V. N., & Persson, B. N. J. (2012). *Effective Viscosity of Conned Hydrocarbons*. Abstract from "Friction: From elementary mechanisms to macroscopic behavior", Berlin, Germany.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Effective Viscosity of Confined Hydrocarbons

Ion M. Sivebaek^{1) 2) 3)*}, Vladimir N. Samoilov^{1) 4)} and Bo N.J. Persson¹⁾

¹⁾ IFF, FZ-Jülich, D-52425 Jülich, Germany

²⁾ Mech. Eng. Dept., Technical University of Denmark, DK-2800 Lyngby, Denmark


³⁾ Novo Nordisk A/S, Research and Development, DK-3400 Hillerød, Denmark

⁴⁾ Moscow MV Lomonosov State Univ., Fac Phys, 117234-Moscow, Russia

*Corresponding author: ims@mek.dtu.dk

We present Molecular Dynamics (MD) friction calculations for confined hydrocarbon films with molecular lengths from 20 to 1400 carbon atoms. We find that the logarithm of the effective viscosity η_{eff} for nanometer-thin films depends linearly on the logarithm of the shear rate γ : $\log_{10} \eta_{eff} = C - n \log_{10} \gamma$, where n varies from 1 (solid-like friction) at very low temperatures to 0 (Newtonian liquid) at very high temperatures, following an inverse sigmoidal curve. Only the shortest chain molecules melt, whereas the longer ones only show a softening in the studied temperature interval $0 < T < 900$ K. The results are important for the frictional properties of very thin (nanometer) films and to estimate their thermal durability. The results are in good accordance with the outcome of Surface Force Apparatus experiments (see e.g. Yamada S. Trib Lett, 13 (3), p 167, 2002)

Figures: Factor n as a function of temperature for three molecule lengths and factor C as a function of n for all investigated systems.



From: Sivebaek, I.M., Samoilov, V.N. and Persson, B.N.J. PRL. 108, 036102 (2012)